# Chancroid Facts

Chancroid is a highly contagious but curable STD caused by a bacteria. It is very common in parts of Asia and Africa and is becoming more common in the U.S.

# Possible Signs and Symptoms

- Women often do not have symptoms.
- Symptoms usually occur within 10 days of exposure
- Chancroid ulcers usually begin as an elevated bump that becomes a puss filled open sore with ragged edges
- Chancres may be tender to the touch—they may be painful in men and unnoticed in women
- Painful lymph nodes may occur in the groin, usually on one side

### Transmission

 Through skin-to-skin contact with an open chancroid ulcer, usually during sexual activity such as vaginal, anal, or oral sex

### Prevention

- Abstain from oral, anal, and vaginal sex
- Consistent, correct use of latex or polyurethane condoms or barriers during oral, anal, or vaginal sex
- Early treatment of STD infection
- If a partner is infected, avoid sexual contact until partner's treatment is completed
- Ensure appropriate treatment of an infected partner

For more information about STDs or about testing and treatment in New Jersey, please call the HIV/STD hotline or visit our web site:

1~800~624~2377

http://www.nj.gov/health/cd/stdhome.htm

### Treatment

- Examination and tests by a health care provider to diagnose and determine best course of treatment
- Chancroid is curable with antibitotics
- Follow-up examination can help ensure that treatment has been successful
- Treatment of sexual partner(s)

# Complications/Effects if Left Untreated

- Increased risk for HIV infection
- Lymph node glands can become infected and painful and require surgery to treat

# Help Stop the Spread of Infection...

- Refrain from sexual contact (oral, anal, or vaginal sex) if you have symptoms
- Seek medical attention, including STD tests
- Refrain from any sexual activity with others until your treatment has been completed
- Take all medications prescribed, even if you no longer have symptoms
- Make sure all recent sex partners are notified if you have chancroid
- Annual STD screenings if you are, or have been, sexually active
- Avoid touching infected areas to avoid spreading chancroid to other parts of your own body



